Understanding sources of predictability at S2S Scale


Andrew W Robertson



Outline

- S2S predictability:
 - Gray zone between weather vs climate predictability
- Phenomena/modes of variability on S2S scale (eg. MJO)
- Slowly varying atmospheric boundary conditions (Soil moisture, snow cover/snow pack, sea ice, SST, ...)

Weather and Climate Forecast Ranges





Atmospheric Predictability

Weather:

Initial Value
Problem
(e.g., baroclinic
waves)

S2S:

Mixed
Initial Value (e.g. MJO)
and Boundary Value
Problem
(e.g. Soil moisture,
snow cover/snow
pack, sea ice, SST)

Seasonal Climate:

Boundary Value Problem (e.g. ENSO SST anomalies)

Daily values

1-10 days

Weekly averages

10-30 days

Monthly or seasonal averages

30-90+ days

TIME AVERAGING!

Predictability of the Second Kind (Lorenz, 1975)



How should the target periods be defined for sub-seasonal forecasts?






FIG. 1. Schematic of the time window and lead time definitions used in this analysis. The horizontal axis represents forecast time from the initial condition. The expression "1d1d" refers to an averaging window of 1 day at a lead time of 1 day. Similarly, "2d2d" represents an averaging window of 2 days at a lead time of 2 days, and so on. Note that 1d1d is what is usually called "day 2" in other papers, and 1w1w is what is usually called "week 2."

Zhu et al (2014, MWR, DOI: 10.1175/MWR-D-13-00222.1)



The Seamless Predictability concept





"The phenomena in the Earth system that occur on all time-scales imply potential predictive power for the atmosphere on that time-scale."



S2S Phase II Research Foci on Sources of Predictability

- MJO prediction and Teleconnections: incl. high impact weather in the tropics/subtropics
- Land Initialization and Configuration: observing system impact on land initialization/S2S forecasts; land/atmosphere processes in S2S models; land surface impacts on extremes
- Ocean and Sea Ice Initialization and Configuration: role of ocean-atmosphere coupling on S2S; sea ice process simulation, initialization, prediction; S2S marine prediction (eg storm surge, fisheries & coral bleaching)
- Stratosphere: role of vertical coupling, stratospheric systematic errors, and impact of quality of stratospheric initial conditions
- Atmospheric Composition: impact of prognostic aerosols on S2S forecasts; level of complexity needed; predictability of aerosols (e.g. dust) & potential forecast value for applications


for Climate and Society

ECMWF Sub-monthly forecast skill

Weekly average CMAP precipitation Anomaly Correlation

Jun-Aug 1992-2008

Li and Robertson (2015)




Skill from
Atmospheric
Initial Conditions

Skill from MJO & Tropospheric Boundary Conditions



ENSO & MJO Precipitation Teleconnections

Niño 3.4 Index




Jun-Aug Anomaly correlation coefficient

• Both MJO and ENSO play a role

MJO RMM Indices

CMAP pentad precip vs RMM: Jun-Aug 1992-2008



International Research Institute

Earth Institute | Columbia University

for Climate and Society

MJO Forecast Horizon now reaches up to 4 weeks ahead






Figure 2. Forecast lead time (in days) when the MJO bivariate correlation between the model ensemble means and control run reaches the nearly of the Theoretical blackulars represent the 95% level of confidence computed from a 10 000 bootstrap re-sampling procedure. [Colour figure can be viewed at wile of the International Diversity of the

MJO Teleconnections (S2S re-forecasts)




Week 3+4 Anomaly Correlation Skill of Teleconnection Indices



Skill mostly in DJF; mostly subseasonal in PNA; interannual in NAO



Stratosphere



Prediction skill of the 1000 hPa Northern Annular Mode for week 3 in the S2S models

For most models, skill is higher following weak vortex conditions. Similar results are found following strong vortex conditions.

SPARC-SNAP



Summary

- Multiple sources of predictability on the 2 weeks to a season range
- Phenomena/modes of variability: Blocking, MJO, SSWs, NAO, ENSO, QBO, ...
- Slowly varying atmospheric boundary conditions: soil moisture, snow cover/snow pack, sea ice, SST, ...
- S2S Phase II will target Land, Ocean, Stratosphere, Aerosols

