

ANNUAL WWW TECHNICAL PROGRESS REPORT ON THE GDPS

COUNTRY: **Australia**

CENTRE: **NMOC Melbourne**

January 2003

1. SUMMMARY OF HIGHLIGHTS (2002):

Meteorological, Oceanographic and Computer Systems:

- 18 March: Changed to Conventional Standardised hours throughout operational NWP system in NMOC.
- 17 April: MOF upgraded (including re-derivation of all coefficients)
- 30 May: BAM3 Physics, featuring unified and optimized code, operationally implemented in GASP.
- 24 July: Convection improvements implemented in GASP.
- 27 August: Sea State Forecast System upgraded (including assimilation of ERS-2 altimeter data and improvements in resolution)
- 3 September: Operational implementation of 1DVAR in LAPS and extension of forecast range to +72 hours.
- 9 September: National EER test for FMD.
- 26 September: Australian region MSLP PAOBs for LAPS generated directly from Horace on-screen analysis.
- 30 September: Southern hemisphere MSLP PAOBs for GASP generated directly from Horace on-screen analysis.
- 28 October: Stopped using bogus GMS moisture data in LAPS and TLAPS.
- 30 October: TOV3 high latitude observational data used in Ozone analysis.
- 6 November: Operational implementation of Coupled Ocean-Atmosphere model (POAMA).
- 25 November: Early cut-off, short forecast, runs of GASP introduced into operations.
- 11 December: TLAPS upgraded and including an extension of forecast range to +72 hours.

2. EQUIPMENT IN USE AT CENTRE:

The following table shows the main computer systems used in NMOC Melbourne, with their basic functions:

FUNCTION	COMPUTER	NO. OF CPUS	MEMORY	OPERATING SYSTEM	DISK STORAGE
Supercomputing (Assimilation and Prediction)	Dual-Node NEC SX-5	Node 1: 16 Node 2: 16	Node 1: 128 GB Node 2: 96 GB	SUPER-UX 10.2 SX-5	2.3 TB
Communications / Message Switching	(i) HP RP5470 (ii) HP RP5470	(i) 2 (ii) 2	(i) 4 GB (ii) 4 GB	(i) HP-UX 11.11 (ii) HP-UX 11.11	200 GB
Scheduling / Data Base / Satellite Post-Processing	(i) HP RP7410 (ii) HP RP7410	(i) 6 (ii) 6	(i) 6 GB (ii) 6 GB	(i) HP-UX 11.11 (ii) HP-UX 11.11	(I) 450 GB (ii) 300 GB
Post-Processing / Graphics Processing	(i) HP D390 (ii) HP D390	(i) 2 (ii) 2	(i) 1500 MB (ii) 1500 MB	(i) HP-UX 10.20 (ii) HP-UX 10.20	(i) 8 GB (ii) 8 GB
Horace	(i) HP C3600 (ii) HP C3600 (iii) HP C3600	(i) 1 (ii) 1 (iii) 1	(i) 512 MB (ii) 512 MB (iii) 512 MB	(i) HP-UX 10.20 (ii) HP-UX 10.20 (iii) HP-UX 10.20	(i) 27 GB (ii) 35 GB (iii) 30 GB
RADAR and Visualisation	(i) SGI Octane (ii) SGI-O2	(i) 1 (ii) 1	(i) 512 MB (ii) 128 MB	(i) IRIX 6.5 (ii) IRIX 6.5	(i) 17 GB (ii) 8.5 GB
web / ftp	4-Node HP lp222r Cluster	2 / Node	2 GB / Node	Linux RedHat 7.3	120 GB / Node
MARS	4-Node IBM RS/6000 SP	High Node 1: 4 High Node 2: 4 Wide Node 1: 2 Wide Node 2: 2	1 GB / cpu	AIX 4.3	1.5 TB
Development	HP RP8400	8	8 GB	HP-UX 11.11	1.8 TB

Peripheral Equipment:

Magnetic Cartridge Archive System:

StorageTek Mass Store 9310 ACS (Automatic Cartridge System)

1 StorageTek 9710 Library Storage Module (with a 37 TB capacity)

6 DLT (Digital Linear Tape) Units

24 drives, for 9840 fibre channel tapes, in silo (8 for SAM-FS, 8 for MARS and 6 for backup)

3 drives for 9940B tapes (for SAM-FS)

Hardcopy Printers/Plotters:

HP DesignJet 1055cm plus and 650C plotters

HP LaserJet 5 SiMX printers

HP LaserJet 4500DN

HP DeskJet 1600CM

Software in use at Centre:

The overall computer environment in the NMOC is mainly Unix. A real-time data base, currently using ORACLE 8.1.7.0.0, is used for storage of observational data and grids from the various NWP systems. The U.S. Navy's Environmental Operational Nowcasting System (NEONS) software is commonly used for accessing the data base. The operational NWP models are written mainly in Fortran, with many of the associated files having the NetCDF structure. Most displays are produced using the NCAR graphics package and IDL. The UK Met. Office's Horace (V3.71) system is used to prepare, in an on-screen mode, manual-computer products for: significant weather prognoses, Australian Region surface analyses and prognoses and Southern Hemisphere surface analyses. Sam-fs is currently being used for magnetic cartridge archives in the NMOC. The Meteorological Archive and Retrieval System (MARS from the ECMWF) is being increasingly used for both research and operations.

Other Systems in use at Centre:

The DIFACS system is used to disseminate a selection of basic analysis and prognostic charts, and some satellite imagery, to the Bureau's regional offices and some outside users. MCIDAS is used for comprehensive interaction and display of satellite imagery and products, observational and gridded data, and is also a major component of the Australian Integrated Forecast System (AIFS). Products from the NWP systems are written to internal and external (www.bom.gov.au) web servers. Magnetic cartridge archives are kept of various NWP products with Australian region analyses available back to 1970 and Southern Hemisphere analyses back to 1972. Hard copy and microfilm archives of charts also exist. An aviation system, which interacts with the WAFS data, is used to view and display the data and prepare the various flight and route forecasts. Regular statistics (including S_1 skill scores, root mean square errors and anomaly correlations), monitoring and comparing the performance of the NMOC's NWP systems (and also some NWP models from overseas centre), are also produced.

3. DATA AND PRODUCTS FROM GTS IN USE:

The following table gives a list of the major observation report types used in the NMOC Melbourne and the approximate numbers received in a 24-hour period:

Observational Data Type or Report Type	Approximate number received during 24 hour period (unless otherwise stated)
SYNOP	41000
SHIP	6400
TEMP	1200
PILOT	820
BUOY	9900
AIREP / AMDAR	20800
SATOB	277000
SATEM	24600
ATOVS (120KM)	115000
METAR	14700
CLIMAT	1500 / month
SATOB_SST	4400
BATHY	140
WAVEOB	900
TRACKOB	620
ACAR	80000
AMV	550000
TOVS1C	2300000
SAT_ALT	53000
QUIKSCAT	1100000

The following Gridded Products are also received in NMOC Melbourne:

- GRIB (ECMWF)
- GRIB (EGRR)
- GRIB (KWBC)
- GRIB (JMA)
- GRID (ECMWF)

4. DATA INPUT STREAM:

Automated. (Some manual intervention is available for correction of reports.) The observational

data, along with NWP gridded data, is stored in a real-time relational data base system (ORACLE/NEONS). An increasing proportion of operationally-produced NWP gridded data is also being stored in MARS.

5. QUALITY CONTROL SYSTEM:


Validity checks are currently confined to within the respective assimilation or analysis schemes. Some gross checking outside these schemes may eventually be installed.

6. MONITORING OF THE OBSERVING SYSTEM:

Monitoring of the observing system is carried out. The quantity of data available is monitored in real-time to ensure that reports are being received reliably and are passed on to the operational systems. For the global system, statistics on the difference between observations and the first guess and analysis fields are routinely prepared to identify any problems with either the analysis system or individual data types. Lists of rejected data are also used to identify unreliable reporting from particular observing platforms.


7. FORECASTING SYSTEM:

There are three major operational analysis and forecast systems (viz. the global GASP, regional LAPS and the tropical TLAPS) in the NMOC Melbourne. A mesoscale version of LAPS, called MESO_LAPS, provides additional high resolution forecast products over 3 smaller domains (viz., Australia, Sydney and Melbourne). The domains for each of these systems are shown in the figure below. The regional and tropical systems are dependent on the global system for their lateral boundary conditions, whereas the 3 MESO_LAPS systems are all nested in LAPS. An additional system, called TC_LAPS, is run to provide tropical cyclone, and other tropical guidance, for the region. (The possible lateral extent of this guidance is also depicted in the figure below.) Manual intervention is used for mean sea level pressure in both the global and regional systems. The resulting hemispheric "pseudo-observations" for mean sea level pressure are disseminated on the GTS. The tropical and Australian region limited area systems both use a tropical cyclone synthetic specification scheme. An additional feature of the tropical system is its dynamical nudging. Output from the global system is also used in the cold start procedure for the Australian region and tropical systems. It is noted that the MESO_LAPS systems do not have their own separate analyses but currently use initial (and boundary) conditions derived directly from LAPS.



Domains of the operational NWP systems in NMOC Melbourne.

The global, Australian region and mesoscale streams have associated sea-state systems. There are a large number of other parts to the basic scheme. These include systems for sea surface temperature analysis, environmental emergency response, generation of weather elements from model output (MOF), amendment and dissemination of aviation products, MCIDAS, archives, verification, display and dissemination of products. A schematic representation of the overall system is shown in the following figure.



Schematic representation of the operational analysis and prediction system in NMO.

7.1 System Run Schedule and Forecast Ranges:

At the present time, the centre produces major analyses at 00 and 12 UTC daily for the globe, Australian region and tropical domains. Global forecasts out to 10 days, Australian region and tropical forecasts out to 72 hours, mesoscale forecasts out to 36 hours, and special tropical cyclone forecasts out to 72 hours are produced off these major analyses. The ECMWF's Supervisor Monitor Scheduler is used to integrate the major part of the operational system and to initiate and monitor the various tasks in the operational NWP suite. An approximate daily schedule for the main operational numerical systems is shown in the table below (with the times during the daylight saving months, November to March, shown in brackets).

System Run Schedule and Forecast Ranges

SYSTEM	BASE TIME (UTC)	APPROXIMATE START TIME (UTC)	FORECAST AVAILABILITY (UTC)	FORECAST RANGE FROM BASE DATE/TIME (HRS)
SSTANAL (REGIONAL)	1200	0115	0130	0
LAPS_PT375	0000	0145 (0045)	0215 (0115)	+72
MESO_LAPS_PT125	0000	0215 (0115)	0335 (0235)	+36
MESO_LAPS_PT050(ME)	0000	0215 (0115)	0245 (0145)	+36
MESO_LAPS_PT050(SY)	0000	0245 (0145)	0330 (0230)	+36
EER PREP (MESO_LAPS_PT050 ME)	0000	0245 (0145)	0300 (0200)	+36
EER PREP (LAPS)	0000	0315 (0215)	0330 (0230)	+48
WAVES (REGIONAL)	0000	0315 (0215)	0325 (0225)	+36
EER PREP (MESO_LAPS_PT050 SY)	0000	0330 (0230)	0345 (0245)	+36
EER PREP (MESO_LAPS_PT125)	0000	0335 (0235)	0350 (0250)	+36
TLAPS375	0000	0345	0420	+72
WAVES (MESO_LAPS_PT125)	0000	0400 (0300)	0430 (0330)	+36
TC_LAPS	0000	0400	0410	+72
EER PREP (TLAPS375)	0000	0500	0520	+48
GASP	0000	0630 (0530)	0800 (0700)	+240
GASP_EPS	0000	0710 (0610)	0900 (0800)	+240
EER PREP (GASP)	0000	0730 (0630)	0830 (0730)	+72
WAVES (GLOBAL)	0000	0730 (0630)	0750 (0650)	+96
TLAPS375	0600	1045	1130	+6
TC_LAPS	0600	1100	1140	+6
LAPS_PT375	0600	1130 (1030)	1230 (1130)	+6
GASP (EARLY)	0600	1200 (1100)	1220 (1120)	+24
LAPS_PT375	1200	1345 (1245)	1415 (1315)	+72
MESO_LAPS_PT125	1200	1415 (1315)	1535 (1435)	+36
MESO_LAPS_PT050(ME)	1200	1415 (1315)	1445 (1345)	+36
MESO_LAPS_PT050(SY)	1200	1445 (1345)	1530 (1430)	+36

EER PREP (MESO_LAPS_PT050 ME)	1200	1445 (1345)	1500 (1400)	+36
EER PREP (LAPS)	1200	1445 (1345)	1500 (1400)	+48
WAVES (REGIONAL)	1200	1515 (1415)	1525 (1425)	+36
EER PREP (MESO_LAPS_PT050 SY)	1200	1530 (1430)	1545 (1445)	+36
EER PREP (MESO_LAPS_PT125)	1200	1535 (1435)	1550 (1450)	+36
TLAPS375	1200	1545	1620	+72
WAVES (MESO_LAPS_PT125)	1200	1545 (1445)	1615 (1515)	+36
TC_LAPS	1200	1600	1610	+72
EER PREP (TLAPS375)	1200	1700	1720	+48
GASP	1200	1830 (1730)	2000 (1900)	+240
GASP_EPS	1200	1910 (1810)	2100 (2000)	+240
EER PREP (GASP)	1200	1930 (1830)	2030 (1930)	+72
WAVES (GLOBAL)	1200	1930 (1830)	1950 (1850)	+96
TLAPS375	1800	2245	2330	+6
TC_LAPS	1800	2300	2340	+6
LAPS_PT375	1800	2330 (2230)	0030 (2330)	+6
GASP (EARLY)	1800	0000 (2300)	0020 (2320)	+24
OCEAN	(Daily)	1000	1005	0
POAMA	(Daily)	2000	0200	+8 months
SSTANAL (GLOBAL)	(Mid-week)	2315 (Mondays only)	2345 (Mondays only)	0
SST (SUB-SURFACE)	(Monthly)	2215 (Tuesdays only)	2245 (Tuesdays only)	0
<u>AD HOC (TOP PRIORITY):</u>				
EER	ANYTIME	ANYTIME	ANYTIME + 30 mins	+72
TC_LAPS	0000 or 1200	ANYTIME	ANYTIME + 45 mins	+72

7.2 Medium-range Forecasting System (4-10 DAYS):

The acronym **GASP** is given to the **G**lobal **A**ssimilation and **P**rognosis system, which produces medium-range forecast products out to 10 days. Post-processed products from this system are disseminated on the GTS in GRIB form, nationally through "DIFACS" and MCIDAS, and also via Radio-facsimile broadcasts (VWM, VMC and VLM).

7.2.1 Data Assimilation, Objective Analysis and Initialization:

Assimilated data: Mean sea level pressure (surface network, ships, drifting buoys), thickness (radiosondes, satellite retrievals), moisture (dew points, satellite precipitable water), wind (rawinsonde, aircraft, geostationary satellites, constant level balloons), cloud-clear radiances (from NOAA orbiting satellites).

Assimilation cycle, including cut-off time: 6 hourly cycling. H+6 cut-off.

Method of analysis: Multivariate statistical interpolation + univariate O.I. for moisture, one-dimensional variational retrievals (1DVAR).

Analysed variables: Geopotential, wind, moisture.

First guess: 6 hour forecast from previous cycle.

Coverage: Global.

Horizontal resolution: Triangular 239.

Vertical resolution: 29 sigma levels (0.991, 0.975, 0.950, 0.925, 0.900, 0.875, 0.850, 0.800, 0.750, 0.700, 0.633, 0.566, 0.500, 0.433, 0.366, 0.320, 0.290, 0.260, 0.230, 0.200, 0.170, 0.140, 0.110, 0.090, 0.070, 0.050, 0.030, 0.020, 0.010)

Initialization: Incremental non-linear normal mode.

7.2.2 Model:

Basic equations: Spectral primitive equations.

Independent variables: latitude, longitude, σ , t

Dependent variables: $\log p^*$, T, q, vorticity, divergence.

Numerical technique:

horizontal: Spectral.

vertical: Finite difference.

time: Semi-implicit semi-Lagrangian.

Integration domain (in horizontal and vertical): Global, surface to 10 hPa (approx.).

Horizontal and vertical resolution, time step: Triangular 239, 29 sigma levels, 600 sec. (approx.).

Orography, gravity wave drag: Both included.

Horizontal diffusion: Included.

Vertical diffusion: Included.

Planetary boundary layer: Included.

Treatment of sea surface, earth surface and soil: Included.

Radiation: Diurnal cycle, diagnostic clouds, interactive optical properties.

Convection (deep and shallow): Included.

Atmospheric moisture: Included.

Boundaries: Stand alone.

Albedo: Climatology.

SST Analysis: Weekly ($1^0 \times 1^0$).

7.2.3 Numerical Weather Prediction Products:

The following table lists some of the post-processed fields that are available from GASP:

Mean sea level pressure	Surface pressure	Evaporation (over ocean)

Surface temperature	Geopotential height	Snow
Temperature	Mixing ratio	Sub-surface temperature
Topography	Zonal velocity	High level cloud
Meridional velocity	Vorticity	Middle level cloud
Divergence	Stream function	Low level cloud
Velocity potential	Vertical velocity	Outgoing longwave radiation
Radiative tendency	Temperature Tendency	Albedo
Mixing ratio tendency	Vorticity surface flux tendency	Net downward radiation at surface
Divergence surface flux tendency	Zonal stress	Roughness length at surface
Meridional stress	Sensible heat flux	Earth albedo
Latent heat flux	Solar heating tendency	Net solar radiation
Precipitation	Soil moisture (over land)	

7.2.4 Operational Techniques for Applications of NWP Products:

The 10 m wind field from GASP is used to drive a global sea-state model.

7.2.5 Ensemble Prediction System:

An Ensemble Prediction System based on the GASP system is currently being run in test mode in real-time in NMOC Melbourne. The 33-member ensemble system is running at the resolution: T119L19 (ie Triangular 119, 19 vertical levels) and producing forecasts out to 10 days, off 00 and 12 UTC. Singular vectors are used to perturb the initial state derived from GASP. These perturbed states are then used as the initial conditions for each of the ensemble members.

7.3 Short-range Forecasting System (0-72 HRS):

The Australian region Limited Area Prediction System (**LAPS**), the Tropical Limited Area Prediction System (**TLAPS**) and the **MESOScale Limited Area Prediction System (MESO_LAPS)** provide short-range forecasting guidance and products that are disseminated nationally through "DIFACS" and MCIDAS, and also via Radio-facsimile broadcasts (VMC, VMW and VLM). TLAPS and TC_LAPS, for specific tropical cyclone guidance, are run on behalf of RSMC Darwin. Again it is noted that, at present, the 3 MESO_LAPS systems do not have their own assimilation, or analysis, but use an initial starting condition derived directly from LAPS. It is also noted that TC_LAPS runs in 2 basic parts. The first preparatory part produces analyses and prognoses over a large domain, or what is called the Large Scale Environment (LSE), and the second part then generates analyses and forecasts at a higher resolution on a relocatable domain centred on the tropical cyclone (and nested within the LSE part).

7.3.1 Data Assimilation, Objective Analysis and Initialization:

Assimilated data: Surface synop, ship, drifting buoy, radiosonde, rawinsonde, GTS TOVS, locally processed TOVS, GTS and locally derived GMS cloud drift winds, aircraft single level winds,

bogus MSLP ("pseudo-observations"), tropical cyclone synthetic data, cloud-clear radiances (from NOAA orbiting satellites).

Assimilation cycle, including cut-off time: 6 hourly cycling; cut-off: LAPS: H+2 hr, TLAPS: H+4 hr.

Method of analysis: Multivariate statistical interpolation + univariate statistical interpolation for moisture, one-dimensional variational retrievals (1DVAR).

Analysed variables: Geopotential, wind, moisture.

First guess: 6 hour forecast from previous cycle.

Coverage: LAPS: 17.125⁰N-65.0⁰S, 65.0⁰E-184.625⁰E

TLAPS: 44.25⁰N-45.0⁰S, 70.0⁰E-188.25⁰E

TC_LAPS: 56.75⁰N-55.0⁰S, 70.0⁰E-164.75⁰W (LSE), 27.0⁰x27.0⁰ (Relocatable)

Horizontal resolution: LAPS and TLAPS: 0.375⁰

TC_LAPS: 0.75⁰ (LSE) and 0.15⁰ (Relocatable)

Vertical resolution: LAPS, TLAPS: 29 sigma levels (0.9988, 0.9974, 0.9943, 0.9875, 0.9750, 0.9625, 0.9500, 0.9250, 0.9000, 0.8750, 0.8500, 0.8000, 0.7500, 0.7000, 0.6000, 0.5000, 0.4500, 0.4000, 0.3500, 0.3000, 0.2750, 0.2500, 0.2250, 0.2000, 0.1750, 0.1500, 0.1000, 0.0700, 0.0500)

TC_LAPS: 19 sigma levels (0.9910, 0.9750, 0.9500, 0.9000, 0.8500, 0.8000, 0.7500, 0.7000, 0.6000, 0.5000, 0.4000, 0.3500, 0.3000, 0.2500, 0.2000, 0.1500, 0.1000, 0.0700, 0.0500)

Initialization: LAPS and MESO_LAPS: Digital filtering technique.

TLAPS and TC_LAPS: diabatic dynamical nudging scheme incorporating GMS IR imagery. (For Tropical Cyclones: Synthetic vortex specification.)

7.3.2 Model:

Basic equations: Grid primitive equations.

Independent variables: x,y,z,t

Dependent variables: P*,T,q,u,v.

Numerical technique:

horizontal: Finite difference.

vertical: Finite difference.

time: Semi-implicit.

Integration domain (in horizontal and vertical):

LAPS: 17.125⁰N-65.0⁰S, 65.0⁰E-184.625⁰E, surface to 50 hPa (approx.)

TLAPS: 44.25⁰N-45.0⁰S, 70.0⁰E-188.25⁰E, surface to 50 hPa (approx.)

MESO_LAPS:

Australian: 4.875⁰S-55.0⁰S, 95.0⁰E-169.875⁰E, surface to 50 hPa (approx.)

Sydney: 30.05⁰S-38.00⁰S, 147.00⁰E-154.95⁰E, surface to 50 hPa (approx.)

Melbourne: 34.05⁰S-41.00⁰S, 141.00⁰E-150.45⁰E, surface to 50 hPa (approx.)

TC_LAPS:

LSE: 56.75⁰N-55.0⁰S, 70.0⁰E-164.75⁰W, surface to 50 hPa (approx.)

Relocatable: 27.0⁰x27.0⁰, surface to 50 hPa (approx.).

Horizontal and vertical resolution, time step:

LAPS: 0.375⁰, 29 sigma levels, 40 sec

TLAPS: 0.375⁰, 29 sigma levels, 40 sec

MESO_LAPS:

Australian: 0.125⁰, 29 sigma levels, 10 sec

Sydney: 0.05⁰, 29 sigma levels, 5 sec

Melbourne: 0.05⁰, 29 sigma levels, 5 sec
 TC_LAPS:
 LSE: 0.75⁰, 19 sigma levels, 40 sec
 Relocatable: 0.15⁰, 19 sigma levels, 15 sec.

Orography, gravity wave drag: Included.
Horizontal diffusion: Included.
Vertical diffusion: Included.
Planetary boundary layer: Included.
Treatment of sea surface, earth surface and soil: Included.
Soil moisture analysis: Included in LAPS, TLAPS, TC_LAPS and MESO_LAPS.
Radiation: Diurnal cycle, diagnostic clouds, interactive optical properties.
Convection (deep and shallow): Included.
Atmospheric moisture: Included.
Boundaries: LAPS, TLAPS and TC_LAPS (LSE): Lateral boundaries from GASP.
 MESO_LAPS (3 domains): Lateral boundaries from LAPS.
 TC_LAPS (Relocatable): Lateral boundaries from LSE TC_LAPS.
Albedo: Climatology.
SST Analysis: Weekly (1⁰x1⁰) - in LAPS, MESO_LAPS (Australian), TLAPS and TC_LAPS.
 Daily (0.25⁰x0.25⁰) - in MESO_LAPS (Melbourne and Sydney).

7.3.3 Numerical Weather Prediction Products:

The following table lists some of the post-processed fields that are available from the Limited Area Prediction Systems:

Mean sea level pressure	Dew point depression	V-component of wind
Surface temperature	Equivalent potential temperature	Potential temperature
Convective precipitation	Ucomponent ageostrophic wind	Divergence
Thickness (1000-500 hPa)	Shearing deformation	Relative humidity
Temperature gradient at 950 hPa	Ucomponent of deformation	Temperature advection
Lifting condensation level	Q vector in x-direction	Stretching deformation
Grassland fire danger index	U-component stream function	Pressure of cloud top
Surface mixing ratio	Moisture advection	Vertical velocity
Boundary layer convective energy flux	Surface pressure	Wet bulb potential temperature
Sensible heat	Non-convective precipitation	Frontogenesis function
U-component of wind	Total precipitation	Vcomponent ageostrophic wind
Temperature	Total totals index	1000-500 hPa shear
Geopotential height	Topography	Vcomponent of deformation
Dew point temperature	Lifting index	Q vector in y-direction
Wind speed	Forest fire danger index	V-component stream function
Vorticity	Mixing ratio	Moisture convergence

Cloud cover	Stream function	
Vorticity advection	Latent heat	

7.3.4 Operational Techniques for Applications of NWP Products:

The 10 m wind field from LAPS and MESO_LAPS are used to drive sea-state models for the Australian region and smaller domains. The Model Output Forecast (MOF) system, which is driven by LAPS and GASP, is used to produce numerous weather elements including temperatures (minimum, maximum, dry bulb, dew point, wet bulb and ground minimum), wind (speed and direction), precipitation, cloud amount, evaporation, sunshine and visibility. Meteograms, giving time series of meteorological variables, are also available. In addition, there is available thunderstorm, based on a decision tree approach, and cold-season tornado (“COLDIES”) guidance, with input from (the Australian domain) MESO_LAPS.

7.4 Specialized Forecasts:

An analogue/regression tropical cyclone model ("TOPEND") is also available for providing guidance to the Australian Tropical Cyclone Warning Centres, but is rarely used. Specialized sea-state forecasts are provided for the North West Cape and Bass Strait gas and oil fields.

7.4.1 Sea Wave Models:

The following table summarises the characteristics of the seastate system in the NMOC:

Domain	Global	Australian Region	Australian MESO_LAPS Region
Numerical Scheme	Deep water 3rd generation (WAM)	Deep water 3rd generation (WAM)	Shallow water 3rd generation (WAM)
Wind data source	GASP (10m)	LAPS_PT375 (10m)	MESO_LAPS_PT125 (10m)
Grid	Latitude/ longitude	Latitude/ longitude	Latitude/ longitude
Resolution	1 ^o	0.5 ^o	0.125 ^o
Nesting	Stand alone	Within Global	Within Regional
Start time of forecast	00,12 UTC	00, 12 UTC	00, 12 UTC
Forecast Period	+96 hrs	+36 hrs	+36 hrs
Initial state	12 hr hindcasting and assimilation of satellite (ERS) altimeter data		No assimilation
Model output	Wind and swell significant wave height, period and direction. Significant wave spectra and probabilities.		
Verification	With respect to rigs, buoys and satellite (ERS) altimeter data		

7.4.2 Storm Surge System:

A system capable of forecasting storm surges caused by tropical cyclones is currently driven by wind and pressure field distributions defined according to the central pressure and maximum wind radius bogussed for tropical cyclones by Darwin RFC.

7.4.3 Air Dispersion Model:

An air dispersion, or transport, system is available for running on demand and can produce forecast trajectories, concentrations (or exposures) and depositions for nuclear accident, volcanic ash, smoke, air-borne virus and other episodes. Currently, the operational Environmental Emergency Response system consists of the Hybrid Single_Particle Lagrangian Integrated Trajectories (HY-SPLIT using Versions 4.0 to 4.6) system, developed at the NOAA Air Resources Laboratory, with meteorological input from the operational NWP systems in NMOC (viz., GASP, LAPS, TLAPS, and MESO_LAPS). Analysed backward trajectories, and retroplumes, are also available from GASP, LAPS and TLAPS in operations.

7.4.4 Solar Ultraviolet (UV) Radiation Forecast System:

Forecasts out to 36 hour forecasts of a UV index (defined as the product of the UV irradiance and a human skin response function) are produced operationally. The system analyses ozone concentrations, available from the global 120 km GTS TOVS data, using 2-dimensional univariate statistical interpolation. Forecasts of the ozone distribution are then computed using isentropic fields derived from GASP output. From vertical profiles of temperature and ozone, the UV index is calculated. Currently, forecasts of the UV index (defined according to the WMO standard categories) are produced, once per day, after completion of the 1200 UTC run of GASP.

7.5 Extended-range Forecasts (10 - 30 DAYS): Not applicable yet.

7.6 Long-range Forecasts (30 DAYS - 2 YEARS):

A three-month rainfall seasonal climate outlook is prepared. Each month, a risk-assessment for three-month total rainfall across Australia is issued mid-month for the three-month period starting the following month. Probabilities are calculated for the three-month total rainfall being in the lowest one-third of historical falls (tercile 1), the middle one-third (tercile 2), and the upper one-third (tercile 3). The technique used is discriminant analysis, with the inputs being derived from recent Sea Surface Temperature (SST) patterns. Subsidiary techniques involved in the forecast model include principal component analysis of SSTs for the Pacific Ocean, Indian Ocean and Southern Ocean, and principal component analysis of rainfall patterns across Australia. SST EOF (Empirical Orthogonal Function) loadings, at one and three months lag, for the Pacific Ocean ENSO pattern and the Indian Ocean pattern are the current predictor inputs. The tercile probabilities, computed across Australia on a $1^{\circ} \times 1^{\circ}$ grid are published in the form of contoured maps, tabulated averages for the 107 Australian rainfall districts, and tabulated interpolations for cities and towns around Australia. Similar outlooks are now provided for above/below median seasonal rainfall and additionally for both maximum and minimum (three-month average) temperatures.

Additional guidance at the rainfall district level is presented in the form of stratified rainfall climatologies based on recent values of the SOI (Southern Oscillation Index). Rainfall outcomes for eastern Australia, obtained from SOI analogues, are also described.

An ocean model for climate monitoring and seasonal prediction is run daily with forcing provided by GASP. Assimilation of ocean (sub-surface) observations is performed every three days.

7.6.1 Ocean-Atmosphere Coupled Model:

The same ocean model, referred to above, is also used to provide the ocean initial conditions and ocean component of a seasonal to interannual coupled ocean/atmosphere general circulation model known as POAMA (Predictive Ocean Atmosphere Model for Australia). Atmospheric initial conditions are obtained from the operational runs of GASP. The atmospheric component of the coupled system is essentially the same model as the operational GASP but run at the reduced resolution: T47L17 (ie Triangular 47, 17 vertical levels). Eight month forecasts are produced daily from this system and used as monthly, or last 30 day, ensemble forecasts for the seasonal outlooks.

8. VERIFICATIONS OF PROGNOSTIC PRODUCTS:

An annual summary of verification statistics for 2002, for the Australian regional and global schemes, is given in the following tables.

The following abbreviations have been used in the tables:

S1 - 30 day mean Teweles skill score over Australian verification area

r - anomaly correlation coefficient between forecast heights and climatology over Australian verification area

PERSIS - 24, 36 or 48 hour persistence prediction

REGN - Australian Regional LAPS (Limited Area Prediction System)

GASP - Global Assimilation and Prognosis model

RMSE - Root Mean Square Error

Note:

The LAPS and GASP results are with respect to their own analyses.

8.1 Skill Scores and Anomaly Correlation Coefficients:

24 HR VERIFICATION STATISTICS - AUSTRALIAN REGION 2002

BASE TIME: 0000UTC - VALID TIME: 0000UTC

			JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
M S L P	S1	PERSIS	57	51	52	54	52	54	53	56	56	65	57	52
	S1	REGN	22	22	21	22	21	19	19	21	19	20	20	20
	r	REGN	.97	.97	.97	.96	.97	.98	.97	.96	.98	.97	.96	.98
5 h o e i h g	S1	PERSIS	48	48	47	45	43	47	45	45	44	47	47	44
	S1	REGN	15	16	15	17	14	13	12	14	12	12	13	14

p a h t	r	REGN	.98	.98	.98	.97	.98	.98	.97	.97	.98	.98	.98	.99
2 5 0 h p a	S1	PERSIS	44	40	44	40	39	42	38	38	36	40	44	42
	S1	REGN	12	12	13	14	12	12	10	10	9	10	12	12
	r	REGN	.98	.98	.98	.97	.98	.98	.97	.97	.98	.98	.97	.99

36 HR VERIFICATION STATISTICS - AUSTRALIAN REGION 2002
BASE TIME: 1200UTC - VALID TIME: 0000UTC

			JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
M S L P	S1	PERSIS	68	63	64	66	63	68	64	69	63	77	66	62
	S1	GASP	28	26	24	24	24	21	23	24	22	22	26	25
	r	GASP	.95	.96	.96	.96	.95	.97	.95	.95	.96	.96	.94	.96
5 0 0 h p a	S1	PERSIS	59	59	59	57	52	56	54	58	50	55	54	54
	S1	GASP	19	19	18	19	17	15	16	18	14	15	17	17
	r	GASP	.97	.97	.97	.97	.97	.97	.96	.96	.97	.97	.96	.98
2 5 0 h p a	S1	PERSIS	54	49	55	49	47	50	46	47	42	47	51	51
	S1	GASP	15	15	16	16	15	13	12	12	11	12	15	15
	r	GASP	.97	.97	.97	.97	.97	.98	.96	.96	.97	.97	.96	.98

48 HR VERIFICATION STATISTICS - AUSTRALIAN REGION 2002
BASE TIME: 0000UTC - VALID TIME: 0000UTC

			JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
M S L P	S1	PERSIS	74	72	72	73	66	71	65	75	67	83	71	67
	S1	GASP	32	31	28	29	28	25	28	29	26	28	31	28
	r	GASP	.93	.93	.93	.94	.93	.95	.92	.93	.94	.94	.90	.94
5 0 0 h p a	S1	PERSIS	60	66	68	62	55	59	55	63	55	58	59	58
	S1	GASP	22	22	22	24	20	18	20	22	18	19	21	20
	r	GASP	.96	.95	.95	.95	.96	.96	.93	.93	.95	.94	.94	.97
2 5 0 h p a	S1	PERSIS	55	54	61	53	49	52	46	50	46	49	55	55
	S1	GASP	19	18	20	20	17	16	15	15	14	16	18	18
	r	GASP	.95	.95	.95	.95	.95	.96	.92	.95	.95	.95	.93	.96

8.2 Root Mean Square Errors (Annual 2002):

(a) Verification against analyses:

AREA	FIELD	RMSE or RMSVE UNIT	+24HR		+72HR		+120HR	
			00UTC	12UTC	00UTC	12UTC	00UTC	12UTC
Northern Hemisphere	500 hPa Height	m	15.2	15.1	41.3	41.2	68.7	68.7
	250 hPa Wind	m/s	5.6	5.6	11.6	11.6	16.7	16.6
Southern Hemisphere	500 hPa Height	m	20.2	20.0	52.2	51.8	81.6	81.2
	250 hPa Wind	m/s	6.2	6.1	13.0	12.9	18.4	18.3
Tropics	850 hPa Wind	m/s	3.0	3.0	4.7	4.6	5.3	5.3
	250 hPa Wind	m/s	5.6	5.6	8.9	8.9	10.4	10.4

(b) Verification against radiosondes:

AREA	FIELD	RMSE or RMSVE UNIT	+24HR		+72HR		+120HR	
			00UTC	12UTC	00UTC	12UTC	00UTC	12UTC
North America	500 hPa Height	m	17.1	16.9	41.1	41.7	67.7	67.0
	250 hPa Wind	m/s	8.3	8.1	14.4	14.1	19.6	19.1
Europe	500 hPa Height	m	18.8	17.4	42.6	41.3	75.9	74.2
	250 hPa Wind	m/s	7.6	7.4	13.5	13.4	19.9	19.8
Asia	500 hPa Height	m	19.0	19.2	39.0	40.4	57.0	58.5
	250 hPa Wind	m/s	7.9	8.2	13.1	13.4	16.5	16.8
Australia/NZ	500 hPa Height	m	17.1	17.1	33.6	34.9	53.1	56.4
	250 hPa Wind	m/s	7.7	7.9	12.8	13.0	17.6	17.6
Tropics	850 hPa Wind	m/s	4.7	4.7	5.7	5.7	6.3	6.3
	250 hPa Wind	m/s	7.0	7.0	8.9	9.2	10.2	10.5
Northern	500 hPa	m	19.4	19.2	44.0	44.2	72.0	71.9

Hemisphere	Height							
	250 hPa Wind	m/s	7.9	7.8	13.5	13.4	18.6	18.5
Southern Hemisphere	500 hPa Height	m	22.1	22.7	43.5	45.2	65.5	69.0
	250 hPa Wind	m/s	8.4	8.8	13.9	14.1	19.0	19.0

9. PLANS FOR THE FUTURE:

Plans for future operational systems in the NMOC Melbourne include:

- . the move and conversion of systems across to a new Central Computing Facility located in a new building (along with all HO staff) in 2004;
- . an ongoing improvement and generalization of data processing and graphical display (including the further use of MARS and METVIEW and, eventually, a GIS);
- . improvement in data monitoring systems;
- . ongoing improvements to LAPS including: full warm running, improvements to cloud and radiation parameterisations, a semi-implicit semi-Lagrangian formulation, more vertical levels, and a non-hydrostatic formulation;
- . the extension of the forecast range for all MESO_LAPS configurations out to +48 hours;
- . the addition of 2 new domains for MESO_LAPS_PT050;
- . the introduction of a LAPS ensemble prediction system;
- . the introduction of a mesoscale assimilation system;
- . enhancements to GASP incorporating: locally derived cloud drift winds and scatterometer winds, increased vertical resolution (50 levels up to 0.1 hPa), 3DVAR;
- . the introduction of a rainfall verification scheme;
- . a move to drive the storm surge model using meteorological input from TC_LAPS;
- . an ongoing improvement to the operational Environmental Emergency Response System including the use of a unified wind input system and an extension to include a number of chemical transformations;
- . the implementation of a gaussian-type plume model for the microscale;
- . the implementation of an urban air quality forecasting system;
- . the further development of the air-borne virus facility within the EER system and, eventually, its incorporation within the microscale dispersion and air quality systems;
- . further development of products from the coupled ocean-atmospheric model for seasonal analysis and prediction.

10. REFERENCES:

Bourke W., Hart T., Steinle P., Seaman R., Embery G., Naughton M. and Rikus L. : "Evolution of the Bureau of Meteorology's Global Data Assimilation and Prediction System. Part 2: Resolution Enhancements and Case Studies." Australian Meteorological Magazine, 44, 19-40, 1995.

Davidson N.E. and Puri K. : "Tropical prediction using dynamical nudging, satellite defined convective heat source and a cyclone bogus." Monthly Weather Review, 120, 2501-2522, 1992.

Davidson N.E. and Weber H.C.: "The BMRC high resolution tropical cyclone prediction system TC-LAPS." *Monthly Weather Review*, 128, 1245-1265, 2000.

Draxler R.R. and Hess G.D. : "An overview of the HYSPLIT_4 modelling system for trajectories, dispersion and deposition." *Australian Meteorological Magazine*, 47, 295-308, 1998.

Greenslade D.J.M.: "The assimilation of ERS-2 significant wave height data in the Australian Region." *Journal of Marine Systems*, 28, 141-160, 2001.

Harris B.A. and Kelly G.: "A satellite radiance bias correction scheme for data assimilation." *Quarterly Journal of the Royal Meteorological Society*, 127, 1453-1468, 2001.

Le Marshall J., Mills G., Pescod N., Seecamp R., Puri K.K., Stewart P., Leslie L.M. and Rea A.: "The estimation of high density atmospheric motion vectors and their application to operational numerical weather prediction." *Australian Meteorological Magazine*, 51, 173-180, 2002.

Le Marshall J., Pescod N., Seaman R., Mills G. and Stewart P. : "An Operational System for Generating Cloud Drift Winds in the Australian Region and Their Impact on Numerical Weather Prediction." *Weather and Forecasting*, 9, 361-370, 1994.

Le Marshall J.F., Riley P.A., Rouse B.J., Mills G.A., Wu Z.-J., Stewart P.K. and Smith W.L.: "Real-time assimilation and synoptic application of local TOVS raw radiance observations." *Australian Meteorological Magazine*, 43, 153-166, 1994.

Mills G.A. and Colquhoun J.R.: "Objective Prediction of Severe Thunderstorm Environments: Preliminary Results Linking a Decision Tree with an Operational Regional NWP Model." *Weather and Forecasting*, 13, 1078-1092 1998.

Puri K., Dietachmayer G.S., Mills G.A., Davidson N.E., Bowen R.A. and Logan L.W. : "The new BMRC Limited Area Prediction System, LAPS." *Australian Meteorological Magazine*, 47, 203-223, 1998.

Seaman R., Bourke W., Steinle P., Hart T., Embery G., Naughton M. and Rikus L. : "Evolution of the Bureau of Meteorology's Global Data Assimilation and Prediction System. Part 1: Analysis and Initialisation." *Australian Meteorological Magazine*, 44, 1-18, 1995.

Stewart P.K. : "RSMC Melbourne - Procedures, Standards, Results of Exercises and Ongoing Products." WMO Expert Team on Environmental Emergency Response. Beijing, China, 20-24 September 1999.

Web Reference:

POAMA: <http://www.bom.gov.au/bmrc/ocean/JAFOOS/POAMA/index.htm>