

Annual WWW Technical Progress Report

On the Global Data Processing and Forecasting System 2004

ARGENTINA

National Weather Service

(www.meteofa.mil.ar , www.meteonet.com.ar)

I. Summary of research development and main operational changes

The Specialized Regional Meteorological Center Buenos Aires (CMRE BUENOS AIRES) has been running an operational regional ten levels primitive equations model since April 1998 (ARPE). The ETA SMN, is running daily since September 2002, and in an operational way, since January 2003. Fields obtained from this models are available on the Internet. Major changes in 2003/2004:

June 2003 Forecasts of temperature and height of tropopause level from the ARPE model are added to the product package.

November 2003 The horizontal resolution of the ETA SMN model was incremented from 0,35°x0,35° to 0,25°x0,25°. The ETA SMN model covers from 90°W to 30°W and from 65°S to 18°S. The initial and boundary conditions are from the GFS model (NCEP). The integration period was extended to 123 hours and 132 hours for the 00 UTC and 12 UTC cycles respectively.

December 2003 A new package of visualization of the ETA model products has been implemented. Post processing jobs run on real time.

January 2004 Initial and boundaries conditions from the ARPE model are replaced by the GFS model for the initialization of a Wave model.

March 2004 Internet 2 is used to download data from the NCEP.

May 2004 ETA SMN model code fix was added.

October 2004 Upgrade of computer architecture. From an Origin 2000 with 8 processors of 250 MHz and Irix 6.4 to an Origin 2002 with 4 processors of 500 MHz and 4 processors of 250 MHz and Irix 6.5.

II. Equipment in use at the center for operational tasks

Function	Computer	CPUs/ Processor	Memory	Disk Storage
Communications and Data quality control system	SG CHALLENGE S Series	1 R4400 200MHz	256MB	2 GB system disk External 4 x 4 GB SCSI disk
Arpe Model	SG INDIGO ² IMPACT	1 R10000 175MHz	128MB	2 GB system disk External 2 x 9 GB SCSI disk
ETA SMN / Wave Model	SG ORIGIN 2002	4 R10000 250 MHz 4 R10000 500 MHz	1400MB	9 GB system disk 36 GB External SCSI disk

Other peripheral equipment and systems are used for database purposes. The National Weather Office is operating with two databases, the operative one, Ideafix, and Oracle. While the latter is used at the present time for historical datasets, work is in progress to replace the operative database with Oracle.

III. Quality control System

Synop and Temp data received at the center are permanently checked for formal and consistencies errors. Buoys, Satem and Satob data are only checked for formal errors. Manual corrections are done when necessary. Other validity checks are performed by the model objective analysis.

IV. Research and development in Data Assimilation and Numerical Forecasting

Model ARPE

Implemented operationally the first time by the Bureau of Meteorology of Australia and adapted later for routine forecasting at New Zealand Meteorological Service. It was adapted to our region by the C.I.M.A. Group from the University of Buenos Aires directed by Dr. M. Nuñez.

Equations: primitive hydrostatic equations

Initialization: vertical mode initialization scheme

Solution technique: a semi-implicit time difference scheme

Physical processes: surface fluxes of momentum, heat and moisture, large scale and convective precipitation, surface temperature and diurnal cycle.

Grid Resolution: 150 Km on the horizontal and 10 levels in the vertical.

Coordinate system: sigma coordinate in the vertical

Forecast period: 36 hours

Objective analysis: a successive correction one (Cressman). The analyzed variables are geopotential heights, temperature, humidity and wind components for ten pressure levels (1000, 850, 700, 500, 400, 300, 250, 200, 150 and 100 hPa); temperature, pressure and humidity at surface and tropopause pressure level.

Data assimilation: performed every twelve hours. The first guess field is generally the twelve hours one predicted by the model in the previous run and in case of model divergence, the climatological field for that month.

Data used: SYNOP, TEMP, BUOYS, SATEM, SATOB and GRID

ETA SMN model

The development of this model began en 1972 by Fedor Mesinger and Zaviša Janjic at the University of Belgrade and the Federal Hydrometeorological Institute of Yugoslavia. During the last decades, the major developments and improvements were done at the National Centers of Environmental Prediction (NCEP).

Equations: Primitive hydrostatic equations. Nonhydrostatic version included

Grid: Arakawa E-grid in horizontal, Philips grid in the vertical.

Resolution: 25km on the horizontal and 38 layers on the vertical.

Solution technique: Split-explicit time differencing, Arakawa-type in space.

Coordinate system: rotated spherical coordinates in horizontal; eta (step mountain) coordinate in vertical. Sigma coordinate version of the model is available.

Physical processes: surface fluxes over land and water; land surface schemes; multilayer soil/vegetation/snowpack land surface model; subgrid mixing; cumulus parameterizations; radiation parameterization; grid scale precipitation parameterization.

Data used: AVN from the GFS model obtained at the NCEP ftp server. Data boundaries are updated every 12 hours. Sea surface temperature, ice/snow coverage and snowdepth information is updated daily and are included in the ETA SMN model initial conditions.

A control routine was added to the ETA model code to prevent spurious generation of kinetic turbulence energy and prevent model divergence.

V. Research and development results for application of NWP products

ARPE model

Analysis and 6 hour forecasts of mean sea level pressure and 1000/500hPa thickness, 850hPa geopotential and dew point, 500hPa geopotential and temperature, 500 hPa vertical wind component, 250hPa geopotential and wind speed, tropopause height and temperature are updated in the National Weather Service Intranet network twice a day. Horizontal and vertical interpolations are made to obtain analyzed horizontal wind components and temperature fields every two degrees of latitude and longitude and forecasted fields every six hours at the seven flight levels used in our region are updated twice a day. Outputs in GRID format are disseminated through the GTS (Global Telecommunication System).

Mean and anomaly fields at all levels are obtained, analyzed and stored on monthly basis.

Some fields from this model are used to obtain an analyzed field of precipitation through the Hydro-estimator technique from NOAA using high resolution GOES-12 pixel information.

ETA SMN model

Analysis and 6 hour forecasts of mean sea level pressure and 1000/500hPa thickness; 2m temperature and humidity; 10m winds; 850hPa geopotential and dew point; 500hPa geopotential and temperature; 250hPa geopotential and wind speed; tropopause level in fl heights; freezing level in fl heights; low, medium, high and total cloud coverage; 24 hour accumulated precipitation fields (convective and large scale); meteograms for selected cities (approximately 100), are available on the Intranet network. Some selected fields and meteogram are available in the Internet as well. Forecasts are updated every 12 hours.

A complete set of variables every 3 hours and accumulated ones up to 120 hours of forecast are available for the forecasts office at the National Weather Service through an html based system.

VI. Development in objective interpretation and verification procedures including performance statistics.

Objective verification of forecast products continued during 2004. The ARPE model performs routine verification of the two kinds: grid-to-observation point and area average observation-to-grid point verifications. The former evaluates temperature forecast against raobs observations while the latter evaluates forecasts against analysis for different fields a parameters. The area of coverage is regional and the period considered is from October 2003 to September 2004 (Tables 1-3).

Verifications of maximum and minimum temperature and precipitation of selected cities forecasted by the ETA SMN model are performed on monthly basis (Fig.1). Evaluation of the official forecasts are included. Verification of temperature on selected levels and location is shown on Table 4.

VII. Plans for future research and development activities.

Continually increase the number/ power of processors from the Origin 2002 and reduce this way the integration time required for the ETA SMN model. Increment of model resolution is planned as well.

Adapt a suitable assimilation and objective analysis scheme to the ETA SMN model and eventually replace the ARPE model.

Use the outputs of the ETA SMN model as initial conditions for the Sea Wave model.

Run a nested model from the ETA SMN with a higher resolution and smaller domain centered at the area of most economical interest on the region.

Use of a model for trajectory and dispersion of volcanic ashes in support of the Volcanic Ashes Advisory Center Buenos Aires.

Obtain an analyzed field of precipitation through the Hydro-estimator technique from NOAA using high resolution GOES-12 pixel information and ETA SMN model analysis.

Develop an operational verification system suitable to the ETA SMN model for standard variables such as temperature, geopotential height, wind, humidity on standard levels as well as variables near the surface and precipitation.

VIII. Other items

Eleven years of analyzed fields using the five levels model and six years (1998-2004) of analyzed fields using the ten levels model (including the operational visualization of meteorological fields using the GRADS software) are available in this Center. Analyzed fields from the ETA model are also available since January 2003.

Objective Analysis and Forecast Area director: Dr. H.H. Ciappesoni.
SMN Scientific Group: Lic. L. Rosso, M. Suaya, M. Gatto and R. Valdivieso.

Table 1: Skill score of Teweles (%) for the 24 hour forecast of geopotential height from the ARPE model (years 2003/2004)

Level (hPa)	Cycle (UTC)	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
100	0	52.6	-	48.0	37.8	34.5	33.7	35.4	40.1	39.0	43.5	46.6	39.4	41.0
	12	52.9	-	47.8	38.3	36.1	35.0	37.4	40.7	39.2	45.9	42.1	42.9	41.7
250	0	38.8	-	39.1	32.5	31.5	34.7	34.2	36.6	31.8	32.9	33.8	31.0	34.3
	12	37.0	-	37.0	32.2	33.4	32.1	33.5	36.1	31.5	31.6	29.6	30.0	33.1
500	0	46.0	-	43.0	37.0	36.4	36.5	38.4	39.1	37.6	39.7	37.9	37.7	39.0
	12	43.8	-	39.6	34.5	36.5	35.1	38.4	38.1	34.2	35.6	35.2	35.6	37.0
850	0	60.4	-	55.7	52.3	54.7	50.5	52.3	58.0	51.8	56.5	51.6	52.6	54.2
	12	58.9	-	54.1	50.3	54.2	49.7	51.7	55.5	51.4	54.9	47.7	50.8	52.7
1000	0	65.7	-	63.9	60.0	52.4	56.4	55.9	59.2	57.1	60.5	61.0	62.0	59.5
	12	66.3	-	62.8	58.8	53.1	56.1	54.0	57.7	56.7	60.3	59.2	60.4	58.7

Table 2: RMS error (m/s) for the 24 hour forecast of geostrophic wind speed from the ARPE model (years 2003/2004)

Level (hPa)	Cycle (UTC)	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
100	0	9.0	-	7.9	8.0	7.9	7.7	8.6	9.5	9.9	9.6	10.3	9.5	8.9
	12	9.3	-	7.5	8.0	8.0	7.9	8.9	9.8	9.7	10.6	8.7	9.3	8.9
250	0	8.8	-	8.6	9.5	10.1	10.9	10.8	12.1	10.1	9.2	10.1	9.2	9.9
	12	8.5	-	8.4	9.4	11.0	9.8	10.7	12.1	10.4	9.3	8.7	8.9	9.7
500	0	5.5	-	5.6	6.1	6.1	6.5	7.8	8.0	7.2	6.6	6.7	6.7	6.6
	12	5.3	-	5.1	5.9	6.4	6.5	8.5	8.7	6.5	6.1	6.0	6.1	6.5
850	0	3.9	-	3.7	4.1	4.1	4.3	4.9	5.0	4.6	4.6	4.6	4.3	4.4
	12	3.7	-	3.8	4.0	4.1	4.6	5.2	4.9	4.7	4.6	4.1	4.4	4.4
1000	0	5.6	-	4.5	5.2	4.5	4.8	5.7	5.6	5.6	5.7	6.1	5.9	5.4
	12	5.1	-	4.6	5.1	4.8	5.4	6.0	5.6	5.5	5.5	5.6	5.4	5.3


Table 3. RMS error of temperature forecast from the ARPE model against observation (years 2003/2004). Location EZEIZA: -34.35S -58.29W

Level (hPa)	Forecast (hour)	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
100	12	3.6	1.5	3.6	3.2	2.2	3.4	4.1	4.4	3.4	3.6	3.4	2.8	3.4
	24	3.1	3.6	4.0	3.3	2.8	3.9	5.2	4.7	3.9	3.6	4.6	4.5	4.0
	36	4.2	4.0	4.2	4.1	3.7	3.8	5.6	7.1	4.8	4.5	4.2	4.4	4.7
250	12	3.7	2.8	4.1	3.5	3.9	3.1	4.1	4.3	3.5	4.0	4.0	4.0	3.8
	24	2.7	2.7	3.4	4.2	4.4	3.5	5.6	4.8	4.1	3.9	4.2	4.6	4.1
	36	3.9	3.2	3.3	4.1	4.4	4.6	5.5	4.7	4.3	5.1	4.7	5.1	4.5
500	12	3.8	2.3	2.3	2.7	2.2	2.9	3.2	3.1	3.1	3.3	3.4	3.3	3.0
	24	2.5	1.9	1.9	2.8	2.7	3.0	3.3	3.2	3.3	2.6	2.3	3.0	2.8
	36	3.8	2.5	1.8	3.4	2.6	2.9	3.1	3.1	3.1	2.7	2.7	2.8	2.9
850	12	2.1	2.3	2.5	2.9	2.6	2.5	3.0	2.6	3.6	2.3	2.2	2.6	2.6
	24	2.1	1.6	2.2	2.8	2.2	2.7	2.1	3.2	3.9	2.2	2.4	2.9	2.6
	36	2.7	2.5	2.7	4.0	3.1	2.7	3.2	4.3	4.0	3.3	3.4	2.8	3.3
1000	12	2.7	2.2	2.6	2.6	2.5	2.3	2.2	2.2	2.4	2.7	3.0	3.0	2.5
	24	3.1	2.9	2.0	2.7	2.1	2.3	1.9	3.1	2.6	2.5	2.9	2.9	2.6
	36	3.7	2.9	2.7	4.1	2.4	2.3	2.2	3.7	3.0	3.5	3.0	2.9	3.1

Table 4. RMS error of temperature forecast from the 12 UT cycle of the ETA SMN model against observation (year 2004) . Location EZEIZA: -34.35S -58.29W

Month	Forecast (hour)	Level (hPa)				
		100	250	500	850	1000
Jan	24	1.7	1.2	1.5	1.8	2.7
	48	1.6	1.4	2.1	2.4	2.8
	72	2.1	1.4	2.1	2.0	2.7
	96	2.6	1.7	2.2	2.9	3.8
Feb	24	1.4	1.4	1.2	1.6	2.4
	48	1.6	1.7	1.4	1.8	2.5
	72	2.2	2.0	2.0	2.5	3.3
	96	2.6	2.4	2.3	3.2	4.2
Mar	24	2.4	1.7	1.3	1.6	1.7
	48	2.8	1.7	1.5	1.7	2.1
	72	3.2	2.0	1.7	1.9	2.2
	96	3.7	1.9	1.8	2.4	2.2
Apr	24	1.6	1.6	1.5	1.5	2.4
	48	2.4	1.4	2.0	1.6	2.7
	72	2.6	1.7	1.8	2.4	3.5
	96	2.9	2.4	2.4	3.2	3.4
May	24	1.3	1.7	1.3	1.3	1.6
	48	1.6	2.0	1.7	1.5	1.6
	72	1.7	2.1	2.0	2.1	2.1
	96	2.0	1.8	1.9	2.6	2.3
Jun	24	1.8	1.0	1.4	2.0	2.5
	48	2.1	1.4	1.9	2.3	2.9
	72	2.0	2.2	2.3	2.1	3.2
	96	2.2	1.5	2.5	2.3	3.1
Jul	24	2.0	2.1	1.7	1.6	1.8
	48	2.3	2.5	2.1	2.3	2.2
	72	2.5	2.7	3.1	2.9	2.2
	96	2.8	2.9	2.9	2.4	2.3
Aug	24	2.2	1.4	1.7	1.1	1.9
	48	2.7	1.8	2.6	2.1	2.1
	72	2.7	2.3	3.1	2.5	2.4
	96	3.2	2.3	3.7	3.2	3.8
Sep	24	2.3	1.7	1.5	1.8	2.1
	48	2.4	2.0	1.6	2.7	2.5
	72	2.7	2.4	1.8	3.1	2.6
	96	3.1	2.7	2.3	3.3	3.4
TOTAL	24	2.0	1.6	1.5	1.8	2.2
	48	2.2	1.9	1.9	2.2	2.4
	72	2.5	2.2	2.2	2.6	2.8
	96	2.6	2.0	2.2	2.5	2.9

a)



b)




Fig. 1: Rate of hits of Precipitation (a) and Maximum/Minimum Temperatures forecasts (b) from the ETA SMN Model and the Official forecast issued by the National Weather Service. The former is based on the 12 UTC cycle and the latter on based on the 17.30 local time forecast. The validity of the forecasts (day 1, 2, etc) are from 0 to 24 hour local time for the selected day.