

RSMC Montréal Report of Activities for 2015

Executive Summary

Regional Specialized Meteorological Centre (RSMC) monthly tests comprised the primary activity during 2015. Hypothetical scenarios were run over Canada, the United States, Australia, Republic of Korea, and the Netherlands. Other activities included incremental updates and improvements to the response procedures, software, and to the joint RSMC secure web pages. The latter are the primary means of communicating transport model products to National Meteorological and Hydrological Services (NMHS), and between RSMCs. RSMC Montréal received requests - operational as well as planned - for inverse modelling support from the Provisional Technical Secretariat (PTS) of the Comprehensive Test Ban Treaty Organization (CTBTO) throughout 2015.

1. Introduction

The Canadian Meteorological Centre (Meteorological Service of Canada, Environment Canada) is designated by the World Meteorological Organisation (WMO) as the RSMC Montréal for the provision of atmospheric transport modelling (ATM) in case of an environmental emergency response. The primary regions of responsibility are WMO Regional Associations (RA) III & IV, which encompass Canada, United-States, Mexico, Central and South America. In addition to emergency response, RSMC Montréal contributes global inverse modelling support to the CTBTO verification system.

2. Operational Contact Information

Canadian Meteorological Centre (CMC)
Environment Canada
2121 Trans-Canada Highway
DORVAL, Québec
Canada H9P 1J3

Business contact: Mr. Nils Ek
Tel : 1 514 421 7207
Fax : 1 514 421 4679
Email : Nils.Ek@canada.ca

Operational contact (24 hours): Shift supervisor
Tel : 1 514 421 4635
Fax : 1 514 412 4639

3. Responses and information on dissemination of products

i. Production of CTBTO meteorological bulletins

Work continues to transfer the production of bulletins containing meteorological data from CTBTO atmospheric monitoring stations from the Canadian Meteorological Centre (CMC)

to Zentralanstalt für Meteorologie und Geodynamik (ZAMG) in Austria. These bulletins had been issued by CMC under header SNCN19 CWAO.

In order for these stations to be officially recognized internationally, the WMO requested each member-state that has CTBTO stations on its territory to assign synoptic codes to identify the stations. The transfer of production of bulletins thus requires each CTBTO station to first be assigned a WMO synoptic identifier. Observations from those stations which have a WMO synoptic identifier are now transmitted by ZAMG under header ISAX30 LOWM in BUFR format.

iv. Dissemination of products

Transport model graphical products and joint statements are posted to secure joint web pages. When requested by the International Atomic Energy Agency (IAEA) these products are also faxed to relevant RSMCs and National Meteorological and Hydrological Services (NMHS). For examples of the graphical products, see Annex 4 of **WMO, 2011**. Throughout 2014, monitoring of RSMC mirror web pages continued in an effort to ensure that they remained congruent.

It is the practice at RSMC Montréal to transmit blank charts to all RSMC mirror websites at the start of each response, before transmitting the actual product charts once the response has begun.

In addition to the other RSMCs, the following countries' NMHSs are in our email and / or fax lists:

Antigua and Barbuda, Argentina, Bahamas, Belize, Bolivia, Brazil, Chile, Colombia, Costa Rica, Cuba, Dominican Republic, Ecuador, Guatemala, Guyana, Mexico, Netherlands Antilles and Aruba, Panama, Peru, Suriname, Trinidad and Tobago, Uruguay and Venezuela. Efforts are made to keep the contact list up-to-date.

v. Response to requests from CTBTO-PTS

There were a total of 18 requests from the PTS of the CTBTO in 2015, 7 between January and June, and an additional 11 during a CTBTO-PTS exercise in October -November.

vi. Other responses

RSMC Montréal participated in a special nuclear exercise on 7 October 2015 with the NMHS in Buenos Aires.

4. Routine operations

Monthly Test:

RSMCs Montréal, Washington and Melbourne hold a joint test on the second Thursday of every month. Following interest demonstrated by other RSMCs, the request to start the exercise is now emailed to all RSMCs. In addition, RSMC Montréal participated in quarterly tests initiated by the IAEA. The following table lists scheduled monthly and

quarterly tests in 2015.

Month	Source location	Initiated by	RSMC providing joint statement
January	Lucas Heights, Australia	Melbourne	Melbourne
February	Wolsong, Republic of Korea	IAEA	RA II and VI RSMCs
March	Susquehanna, PA, USA	Washington	Montréal
April	Bruce NPP, ON, Canada	Montréal	Washington
May	Lucas Heights, Australia	IAEA	RA III RSMCs (Melbourne)
June	Dresden NPP, IL, USA	Washington	Montréal
July	Lucas Heights, Australia	Melbourne	Melbourne
August	Borssele, Netherlands	IAEA	RA I and RA VI RSMCs
September	Point Lepreau NPP, NB, Canada	Montréal	Washington
October	Lucas Heights, Australia	Melbourne	Melbourne
November	Nine Mile Point NPP, NY, USA	IAEA	RA III and IV RSMCs (Washington)
December	Monticello NPP, MN, USA	Washington	Montréal

5. Lessons learned and significant operational or technical changes:

Upgrades to the software used to run the operational atmospheric transport and dispersion model were implemented periodically during 2015.

6. Operational issues and challenges:

Faxing of products to NMHSs has been discontinued for monthly tests, as email has become the preferred method of communications and faxes exhibit a high failure rate. Faxes will to be sent to RA III and IV NMHS upon request from the IAEA.

7. Other activities:

In response to Action 16 of WMO-CBS Nuclear ERA meeting of October 2013, RSMC Montreal worked in parallel with RSMC Vienna, to develop some examples of radioactive cloud simulations based upon realistic emissions of a major incident, to investigate the potential development of aviation guidance for such incidents. A report was written jointly with RSMC Vienna, presenting the examples and findings. This report was submitted to the Meeting of the CBS Expert Team on Emergency Response Activities, held in Buenos Aires, Argentina, 30 November – 4 December, 2015. RSMC Montreal participated in tests of the production of Time of Arrival charts, with RSMC Obninsk leading the tests. These were conducted on 2 June 2015 and 27 October 2015. There was some initial difficulty in producing the charts in conformity with the specified standard for the desired time intervals and colours with hatching. The work is ongoing.

8. Summary and status of the operational atmospheric transport and dispersion models:

Current global weather conditions and forecasts are available at CMC at all times, to provide, in real time, the necessary input to the ATM, and for their evaluation and interpretation.

For forecasts, CMC operational uses the Canadian Global Environmental Multiscale (GEM) numerical weather prediction (NWP) model. Three configurations of GEM are available: global, regional, and high resolution. The global GEM has a uniform horizontal resolution (25 km) and is used to run the data assimilation cycle, providing analyses and medium term forecast guidance. The grid spacing of the regional configuration is approximately 10 km over North America. High resolution configurations of the GEM model operate at a resolution of 2.5 km and cover all but the northernmost areas of Canada.

i. The Modèle Lagrangien de Dispersion de Particules d'ordre zéro (MLDP0)

This is a Lagrangian particle dispersion model designed for dispersion problems occurring at regional and global scales and is described in detail in D'Amours and Malo, 2004, and D'Amours et al. 2015.. Dispersion is simulated by calculating the trajectories of a large number of air particles (or parcels). Large scale transport is calculated as the displacement due to the NWP-resolved wind field, while discretized stochastic differential equations account for the unresolved turbulent motions. Turbulent vertical mixing is modelled with a random displacement equation based on a diffusion coefficient. Lateral (horizontal) turbulent diffusion is modelled according to a first order Langevin stochastic equation for the components of the horizontal wind that are not resolved by NWP.

MLDP0 is an off-line model and requires 3-D meteorological fields (wind, moisture, temperature and geopotential heights) from a NWP system. At RSMC Montréal these are obtained from the GEM model forecast and analysis system in Global, Regional, or high resolution configuration.

Dry deposition is modeled using a deposition velocity. The deposition rate is calculated as a proportion of the tracer material carried by particles in a layer adjacent to the ground surface. Wet deposition will occur when a particle is presumed to be in a cloud. The tracer removal rate is proportional to the local cloud fraction.

The source term is controlled through an emission scenario module which allows different release rates of radionuclides over time. MLDP0 can be run for a large number of isotopes (Cs-137 by default) as well as for volcanic ash (D'Amours et al, 2010) or an inert gas tracer.

For volcanic eruptions, a particle size distribution can be used to model the gravitational settling effects in the trajectory calculations according to Stokes' law. The total released mass can be estimated from an empirical formula derived by Sparks *et al.*, 1997, which is a function of particle density, plume height and effective emission duration (Malo, 2007).

In MLDP0, tracer concentrations at a given time and location are calculated by averaging the residence time of the particles, during a given time period, within a given sampling volume, and weighting it according to the amount of material carried by the particles. Concentrations are expected to be estimated more accurately near the source with a Lagrangian model than with an Eulerian model.

MLDP0 operates on a polar stereographic grid and can run in both northern and southern hemispheres. The grid size and resolution define the geographical domain. More than 30 horizontal grids are now available, some of which are listed below:

- 50 km (687×687), (477×477), (400×400) and (334×334)
- 33 km (722×722), (606×606), (505×505), (400×400) and (229×229)
- 15 km (503×503) and (251×251)
- 10 km (229×229)
- 5 km (457×457)
- 2 km (300×300)

A global configuration also exists at horizontal resolution of 1° (360×181). MLDP0 can be executed in backward (adjoint) mode. The model has been used extensively in this configuration in the context of the WMO-CTBTO collaboration. The vertical discretization is made for 25 levels in the SIGMA, ETA or HYBRID terrain following coordinates depending on the version of the GEM NWP model used.

ii. Trajectory model

This model uses winds directly from the GEM analyses and/or forecast model. The wind fields are available every hour in forecast mode and every 3 hours in diagnostic mode. Initial positions of one or more air parcels in a column are specified, and the parcels are then incrementally displaced, using time and spatial discriminations of the local three-dimensional wind field. It is assumed that air parcels preserve their identity as they are transported in the wind.

The model has been validated using back-trajectories from stations that measured concentrations of tracers from a single source (D'Amours 1998). The back-trajectories converge remarkably well towards the tracer source location. On the other hand, the lack of a boundary layer treatment and the assumption air parcel identity preservation are reflected in the results, which indicate vertical motions that are not in line with the observations.

9. Plans for 2016-2017:

- The schedule of routine monthly tests for all of 2016 has been set up in collaboration with RSMCs Washington and Melbourne. Each RSMC will select the simulated accident location and write the joint statement, on a rotating basis. Quarterly tests are also scheduled with the IAEA.
- Final implementation of an upgraded version of the MLDP Lagrangian transport and dispersion model that will include zero-order, first-order, and backward-adjoint capabilities.
- Test and implement hourly meteorological fields in diagnostic and forecast modelling

References

- D'Amours, R., 1998: Modelling the ETEX plume dispersion with the Canadian Emergency Response Model, *Atmospheric Environment*, **32**, 4335-4331
- D'Amours, R., and Malo, A., 2004, “*A Zeroth Order Lagrangian Particle Dispersion Model: MLDP0*”, Internal report, Canadian Meteorological Centre, Environmental Emergency Response Section, Dorval, Québec, Canada, 18 pp.
- D'Amours, R., A. Malo, R. Servranckx, D. Bensimon, S. Trudel, and J.P. Gauthier-Bilodeau (2010), *Application of the atmospheric Lagrangian particle dispersion model MLDP0 to the 2008 eruptions of Okmok and Kasatochi volcanoes*, J. Geophys. Res., 115, D00L11, <http://dx.doi.org/10.1029/2009JD013602>.
- D'Amours, R., Malo, A., Flesch, T., Wilson, J., Gauthier J.-P., Servranckx, R. (2015), *The Canadian Meteorological Centre's Atmospheric Transport and Dispersion Modelling Suite*, *Atmosphere-Ocean*, **53** (2), 176–199, <http://dx.doi.org/10.1080/07055900.2014.1000260>
- Malo, A., 16 November 2007, “*Total Released Mass Calculation for Volcanic Eruption in CMC's Long-Range Transport and Dispersion Model MLDP0*”, Internal Publication, Canadian Meteorological Centre, Environmental Emergency Response Section, Dorval, Québec, Canada, 2 pp.
- WMO, 2011: Documentation on RSMC Support for Environmental Emergency Response. *WMO-TD/No. 778*. Available online at <http://www.wmo.int/pages/prog/www/DPFSERA/td778.html>